

POSITRON FRACTION IN THE CMSSM*

WIM DE BOER,[†] MARKUS HORN, CHRISTIAN SANDER

*Institut für Experimentelle Kernphysik, University of Karlsruhe, Postfach 6980,
D-76128 Karlsruhe, Germany*

DMITRI KAZAKOV

*Bogoliubov Laboratory of Theoretical Physics, JINR, 41980 Dubna, Moscow
Region, Russian Federation*

A fit to the present cosmic ray positron fraction can be considerably improved, if in addition to the positron production by nuclear interactions in the universe the possible contribution from supersymmetric dark matter annihilation is taken into account. We scan over the complete SUSY parameter space of the Constrained Minimal Supersymmetric Model (CMSSM) and find that in the acceptable regions the neutralino annihilation into $b\bar{b}$ quark pairs is the dominant channel with hard positrons emerging from the semileptonic decays of the B-mesons.

1. Introduction

The cosmic ray positron fraction at momenta above 7 GeV is difficult to describe by the background only hypothesis¹. A contribution from the annihilation of neutralinos, which are the leading candidates to explain the cold dark matter in the universe, can improve the fits considerably^{2,3,4}. The neutralinos are the Lightest Supersymmetric Particles (LSP) in supersymmetric extensions of the Standard Model, which are stable, if R-parity is conserved. This new multiplicative quantum number for the supersymmetric partners of the Standard Model (SM) particles is needed to prevent proton decay and simultaneously prevents the LSP a) to decay into the lighter SM particles and b) can only interact with normal matter by producing additional supersymmetric particles. The cross sections for the latter are typically of the order of the weak cross sections, so the LSP is “neutrino-like”, i.e. it would form halos around the galaxies and consequently, it is

*Preprint:IEKP-KA/2002-16

[†]E-mail: wim.de.boer@cern.ch

an excellent candidate for dark matter.

In addition to being of interest for cosmology, supersymmetry solves also many outstanding problems in particle physics, among them⁵: 1) it provides a unification of the strong and electroweak forces, thus being a prototype theory for a Grand Unified Theory (GUT) 2) it predicts spontaneous electroweak symmetry breaking (EWSB) by radiative corrections through the heavy top quark, thus providing a relation between the GUT scale, the electroweak scale and the top mass, which is perfectly fulfilled 3) it includes gravity 4) it cancels the quadratic divergencies in the Higgs mass in the SM 5) the lightest Higgs mass can be calculated to be below 125 GeV in perfect agreement with electroweak precision data, which prefer indeed a light Higgs mass. Therefore it is interesting to study the positron fraction from neutralino annihilation in the reduced region of SUSY parameter space, where these constraints are satisfied and compare it with data, as will be done in the next section. It should be noted that the positrons from the annihilation of heavy particles leads to a much harder energy spectrum than the positrons expected from nuclear interactions, thus leading to a significant change in the shape of the spectrum. In addition, in the positron fraction, defined as $e^+/(e^- + e^+)$, many systematic uncertainties related to fluctuations in the nuclear densities and propagation in the universe, cancel.

2. Neutralino Annihilation in the CMSSM

In the Constrained Minimal Supersymmetric Model (CMSSM) with supergravity mediated breaking terms all sparticle masses are related by the usual GUT scale boundary conditions of a common mass m_0 for the squarks and sleptons and a common mass $m_{1/2}$ for the gauginos. The parameter space, where all low energy constraints are satisfied, is most easily determined by a global statistical analysis, in which the GUT scale parameters are constrained to the low energy data by a χ^2 minimization. For details we refer to previous publications⁶. The l.h.s. of Fig. 1 shows a typical running from the common masses at the GUT scale to low energies. The squarks and gluinos get a higher mass than the sleptons due to the gluonic contributions from the strong interactions. If $m_{1/2}$ is not strongly above m_0 , the lightest mass is the supersymmetric partner of U(1) gauge boson, the bino, which mixes with the W^3 boson and spin 1/2 higgsinos to neutralinos. The low energy value of the LSP is roughly 0.4 times its starting value at the GUT scale, i.e. $0.4m_{1/2}$. The gaugino fraction of the LSP




Figure 1. Left: The running of the masses between the GUT scale and the electroweak scale for $\tan \beta = 35$. Right: The gaugino fraction as function of m_0 and $m_{1/2}$ for $\tan \beta = 35$. For lower values of $\tan \beta$ the gaugino fraction is even closer to one. The region with gaugino fraction zero is excluded, since the LSP would be charged, which is excluded because the dark matter is neutral.

is defined as $(N_{i,1}^2 + N_{i,2}^2)$, where the coefficients determine the neutralino mixing $\tilde{\chi}_i^0 = N_{i,1}\tilde{B} + N_{i,2}\tilde{W}^3 + N_{i,3}\tilde{H}_1^0 + N_{i,4}\tilde{H}_2^0$. As shown on the r.h.s of Fig. 1 the gaugino fraction is close to one for LSP masses above 100 GeV, i.e. $m_{1/2} > 250$ GeV. The gaugino fraction is important, since the neutralino properties are quite different for a pure gaugino or pure Higgsino.

Neutralino annihilation can occur through Z- and Higgs exchanges in the s-channel and sfermion exchange in the t-channel. This annihilation in the halo of the galaxies will produce antimatter at high momenta, thus anomalies in the spectra of positrons and antiprotons provide an excellent signal for dark matter annihilation. The neutralino is a spin 1/2 Majorano particle, so it obeys the Pauli principle, which implies an asymmetric wave function or spins antiparallel for annihilation at rest. Final states couple most strongly with total spin 1, so light particles are “helicity-suppressed”. The cross section for $b\bar{b}$ grows quickly with $\tan \beta$ and dominates for large $\tan \beta$, as shown in Fig. 2.

Fig. 3 shows the fit to the data for different regions of parameter space with different main annihilation channels. The fits were done with DarkSusy and follow the same principle as the ones from Ref. ², i.e. the background is left free within a normalization error of about 20% and the signal can be enhanced by a boost factor, which takes into account the uncer-




Figure 2. The thermally averaged cross section times velocity for neutralino annihilation as function of m_0 and $m_{1/2}$ for different final states ($b\bar{b}$, $t\bar{t}$, W^+W^- , $\tau^+\tau^-$) at $\tan\beta=50$. The neutralino mass equals $\approx 0.4m_{1/2}$ in the CMSSM, so in the plots the neutralino varies from 40 to 400 GeV along the front axis. The $b\bar{b}$ final state grows with $\approx \tan^2\beta$ and dominates for $\tan\beta$ above 5. Note the different vertical scales.

tainty of the relic density in our galaxy due to possible clumpiness of the dark matter.

It should be noted, that the fits with W^+W^- and $t\bar{t}$ final states are excluded by the present electroweak data, which requires $\tan\beta$ above 5, in which case the $b\bar{b}$ final state dominates. Unfortunately the data are not precise enough to prefer a certain value of the LSP mass. However, the present balloon experiments correspond to only a few days of data taking. With future experiments, like PAMELA⁹ on a russian satellite or AMS-02¹⁰ on the ISS (International Space Station), one will take data for several years, thus being able to decide if the shape of the positron fraction originates




Figure 3. Fits to the data for $\tan\beta = 1.6$ and $\tan\beta = 50$. In the first case W^+W^- is the dominant annihilation channel, but in the second case $b\bar{b}$ dominates. In both cases the χ^2 reduces from 56 for the background only fit for 27 data points to around 31, if neutralino annihilation is included.

from two components: a hard component from neutralino annihilation and a softer component from nuclear interactions. Taking the χ^2 difference between the fit with a single component and two components as indication of neutralino annihilation, one can predict what the data after one year of data taking with AMS-02¹⁰ with an acceptance of more than $0.04 \text{ m}^2 \text{ sr}$ will look like. This is indicated by the points in Fig. 4. Here we assumed a boost factor of 90 as given by the fit to the present data. Clearly, the positron spectra can provide strong evidence for neutralino annihilation and the data will be accurate enough to precisely determine the neutralino mass, as shown by the r.h.s. of Fig. 4. We like to thank Drs. L. Bergström, J. Edsjö and P. Ullio for helpful discussions.

References

1. S. W. Barwick *et al.* [HEAT Collaboration], *Astrophys. J.* **482** (1997) L191 [arXiv:astro-ph/9703192].
M. A. DuVernois *et al.*, *Astrophys. J.* **559** (2001) 296.
J. Alcaraz *et al.* [AMS Collaboration], *Phys. Lett. B* **484** (2000) 10 [Erratum-ibid. B **495** (2000) 440].
R.L. Golden *et al.*, *Astrophys. J.* **457** (1996) L103.
G. Barbiellini *et al.*, *Astron. & Astrophys.* **309** (1996) L15.
2. E. A. Baltz *et al.*, *Phys. Rev. D* **65** (2002) 063511 [arXiv:astro-ph/0109318].
3. G. L. Kane, L. T. Wang and T. T. Wang, *Phys. Lett. B* **536** (2002) 263




Figure 4. Expected statistics after one year of data taking with AMS-02 for a neutralino of 200 GeV and assuming a boost factor of 90 required by the best fit to the present data. On the r.h.s. the χ^2 contour of the CMSSM fit to the Monte Carlo data on the left. Note the precise determination of $m_{1/2}$, which implies a precise determination of the LSP mass. The region labeled “excl. LSP” is excluded because the LSP is the charged stau; a charged stable particle is not a dark matter candidate.

[arXiv:hep-ph/0202156].

J. R. Ellis, J. L. Feng, A. Ferstl, K. T. Matchev and K. A. Olive, arXiv:astro-ph/0110225.

M. Kamionkowski and M. S. Turner, Phys. Rev. D **43** (1991) 1774.

4. W. de Boer, C. Sander, M. Horn and D. Kazakov, Nucl. Phys. Proc. Suppl. **113** (2002) 221; arXiv:astro-ph/0207557.

5. W. de Boer, Prog. Part. Nucl. Phys. **33** (1994) 201 [arXiv:hep-ph/9402266] and references therein.

6. W. de Boer *et al.*, Z. Phys. C **67** (1995) 647 [arXiv:hep-ph/9405342].

W. de Boer *et al.*, Z. Phys. C **71** (1996) 415 [arXiv:hep-ph/9603350].

W. de Boer *et al.*, Phys. Lett. B **438** (1998) 281 [arXiv:hep-ph/9805378].

7. J. Gondolo, J. Edsjö, L. Bergström, P. Ullio, and E.A. Baltz, astro-ph/0012234.

8. A. W. Strong and I. V. Moskalenko, Adv. Space Res. **27** (2001) 717, [arXiv:astro-ph/0101068]; Phys. Rev. D **60** (1999) 063003 [arXiv:astro-ph/9905283].

9. V. Bonvicini *et al.* [PAMELA Collaboration], Nucl. Instrum. Meth. A **461** (2001) 262.

10. J. Alcaraz *et al.* [AMS Collaboration], Nucl. Instrum. Meth. A **478** (2002) 119.